



NIRMALA COLLEGE OF PHARMACY MUVATTUPUZHA



NATIONAL CONFERENCE

Nano-based Drug Delivery Systems; Recent Developments and Future Prospects

7 OCTOBER 2023

ASSOCIATING PARTNERS

INDIAN PHARMACEUTICAL ASSOCIATION



**JOURNAL OF INNOVATIONS IN
APPLIED PHARMACEUTICAL SCIENCES**



**INNOVATION AND
ENTREPRENEURSHIP
DEVELOPMENT CENTRE**



**INSTITUTION'S
INNOVATION
COUNCIL**
(Ministry of HRD Initiative)



HOME / ARCHIVES / Volume-8, Issue-3-5, 2023



National Conference on Nano-based Drug Delivery Systems; Recent Developments and Future Prospects conducted By Nirma College of Pharmacy, Muvattupuzha, in association with Indian Pharmaceutical Association on 7 October 2023

RESEARCH ARTICLE(S)

ASSESSMENT OF PHYSICAL FUNCTIONING IN RHEUMATOID ARTHRITIS PATIENTS AFTER RITUXIMAB THERAPY USING HEALTH ASSESSMENT QUESTIONNAIRE-DISABILITY INDEX

ANNA MARIA JOY , AKSHARA SHAJI , SHANIYA MATHEW , DR.SUJA ABRAHAM

Pages 1-4

[VIEW PDF](#)

TOXICITY PROFILE OF CHEMOTHERAPY REGIMENS FOR MULTIPLE MYELOMA PATIENTS USING CTCAE CRITERIA

ANTONY V R, ARPITH ANTONY, HELAN KURIAN, JEEVA ANN JIJU, TIMY THOMAS, JITHIN SUNNY, SUJA ABRAHAM

Pages 5-7

[VIEW PDF](#)

ISOLATION OF EMBELIN FROM EMBELIARIBES BERRIES FOR THE DEVELOPMENT OF TOPICAL ANTI-INFLAMMATORY PREPARATION

DR. R. BADMANABAN, MARIA S.PADATHIL , HANNA PARVEEN, DONA MERIN JOY, SHAHANA MAJEED, JOYCYMOLS, DR. DHRUBO JYOTI SEN

Pages 8-18

[VIEW PDF](#)

DESIGN AND CHARACTERISATION OF TOPICAL EMULGEL CONTAINING NEEM OIL FOR ITS ANTIDANDRUFF PROPERTIES

EBY GEORGE, DR DHANISH JOSEPH, ABITHA N JABBAR, KHANSA BEEGAM M A, NIMISHA JOSEPH, MAHIMA FRANCIS, ANJU BOBAN, ANN MARIA ALEX

Pages 19-23

[VIEW PDF](#)

DEVELOPMENT OF IMPLANTABLE DRUG DELIVERY SYSTEM OF EMBELIN FOR THE TREATMENT OF BREAST CANCER

RINCY. K. K, DR. DHANISH JOSEPH, BINSHA URUMEEES, ANN MARIYA JOSE, ATHIRA ANILAN

Pages 24-28

[VIEW PDF](#)

COMPARATIVE INSILICO DOCKING STUDY INVOLVING ANTAGONISTIC ACTIVITY OF COUMARINDERIVATIVES ON EGFR AND CDK2

RIYA ANN THOMAS, EVA SARA SUNIL, ANNA ABEL FERNANDEZ, SOORYA ANIL, ANJANA ANTONY, ANN MARIA DAVIS, GODWIN THOMAS, SARANYA T S, GREESHMA SREERAM, DR. ELIZABETH ABRAHAM P

Pages 29-35

[VIEW PDF](#)

ASSESSMENT OF PATIENT KNOWLEDGE, PRACTICE AND ADVERSE EVENTS OF INSULIN ADMINISTRATION AND STORAGE TECHNIQUES IN PATIENTS WITH DIABETES

ANTRIYA ANNIE TOM, NAMITHA ANTONY, PAVITHRA ASHOK, MUHAMMAD ABDUL KHADIR PS, JUHY JOJO

Pages 42-46

[VIEW PDF](#)

FORMULATION AND EVALUATION OF HERBAL AFTERSHAVE GEL

CELU MARIYA FRANCIS, RIYA GEORGE, ANASWARA SANKAR, ANCI I J, MANJU MARIA MATHEWS, BADMANABAN R

Pages 47-50

[VIEW PDF](#)

EVALUATION OF ANTIMICROBIAL ACTIVITY OF A HERBAL MIXTURE

DEEPA JOSE , SINI BABY, SUJJALA SUBASH, GIFTY LAWRENCE, ANEESA ANOOB , LINTA JOSE

Pages 59-63

[VIEW PDF](#)

ONLINE SUBMISSION



Online ISSN:2455-5177

CODEN (CAS-USA): JIAPAW

Impact Factor: 5.832

Journal Archived in



KEYWORDS



CURRENT ISSUE

Volume 8, Issue 3-5, 2023

Volume 8, Issue 3-5, 2023

Volume 8, Issue 3-5, 2023

INFORMATION

[For Readers](#)

[For Authors](#)

[For Librarians](#)

[Flag Counter](#)

EFFICIENT MICROWAVE SYNTHESIS OF COUMARIN DERIVATIVES WITH EVALUATION OF THEIR ANTIOXIDANT AND ANTI-INFLAMMATORY PROPERTIES
ANZIYA P A, SARANYA T S, ANJALI K, ANJALI KRISHNA, SINI BABY, DIVINE P DANIEL Pages 124-130

[VIEW PDF](#)

COSMETIC USE RELATED ADVERSE EVENTS AND NEED FOR COSMETOVIGILANCE
MERRIN JOSEPH, KARISHMA SHAJI, MAHIN T M, NANDANA P B, KRISHNA DAS Pages 64-71

[VIEW PDF](#)

A RETROSPECTIVE STUDY OF CLINICAL PROFILE OF VIPER BITE CASES IN SELECTED HOSPITALS IN CENTRAL KERALA
ANUMOL SAJU, ANTRIYA ANNIE TOM, ABY PAUL, SWAPNA SAJU, DONA JOHNSON, JESYLN JOE THOMAS, KUTTIKADEN Pages 72-74
JOY STEFFI, JOYAL M JOLL

[VIEW PDF](#)

FORMULATION AND EVALUATION OF HERBAL TOOTHPASTE CONTAINING EUPATORIUM TRIPLINERVISLEAF EXTRACT
VIDYA PETER, ROSNA BABU , SHERRY SEBASTIAN, ANGEL JAEMON, ANGEL JAEMON, ANAGHA V T, JEEVAN SAJEEV Pages 36-41

[VIEW PDF](#)

IN VITRO SCREENING OF ICACINACEOUS PLANTS INDIGENOUS TO KERALA
DR.ELIZABETH ABRAHAM P, FRINTO FRANCIS, PRADEEP R NAIR, ATHUL RAJ, RAJI RAJAN, ANAMIKA K. NAIR, PROF.DR.BADMANABAN.R Pages 51-58

[VIEW PDF](#)

FORMULATION AND EVALUATION OF BUCCAL FILM OF AN ANTIHYPERTENSIVE DRUG
ASHINAA BENEDICT, IRIN ROSE PAUL, DR. MANJU MARIA MATHEWS, DR. BADMANABAN R Pages 75-80

[VIEW PDF](#)

A PROSPECTIVE SURVEY TO ASCERTAIN THE SYMPTOMS, HEALTH ISSUES AND SUBSEQUENT OTC MEDICATION USAGE DURING MENSTRUATION AMONG COLLEGE STUDENTS
MINTU GEORGE, ANAGHA MELBIN, MARY PAUL DOMINIC, RESHMA DOMINIC, AYSHA SAJA P.S, JOBIN KUNJUMON Pages 81-84
VILAPURATHU

[VIEW PDF](#)

A CROSS SECTIONAL STUDY TO ANALYSE THE ADR REPORTED IN A HOSPITAL DURING THE PAST THREE YEARS
SANGEETHA SUKUMARAN, VARSHA ELIZABETH JOBY, AMALA JOSEPH, APARNA JESTIN, JITHIN N P, SUMAYYA B Pages 85-89
MUHAMMED, SUNU SEBASTIAN, JOBIN KUNJUMON VILAPURATHU

[VIEW PDF](#)

FORMULATION AND EVALUATION OF PREUNGUAL DELIVERY SYSTEM CONTAINING EUGENOL FOR THE TREATMENT OF ONYCHOMYCOSIS
MINI ELIAS, FLOWERLET MATHEW, GOURISREE T, ANILA RAJAN, ASHLY DAVIS Pages 90-94

[VIEW PDF](#)

FORMULATION AND EVALUATION OF FLOATING CONTROLLED DRUG DELIVERY OF ANTI-ULCER DRUG LOADED MICROBALLOONS
BINDUMOL K C, FLOWERLET MATHEW, SHALOM SUNIL, ANGEL JOSE Pages 95-100

[VIEW PDF](#)

PREPARATION AND EVALUATION OF FLOATING DRUG DELIVERY SYSTEM (FDSD) CONTAINING AN ANTIVIRAL DRUG
TEENA MOHAN, MARIYA SUNNY, MANJU MARIA MATHEWS, BADMANABAN R Pages 105-109

[VIEW PDF](#)

FORMULATION AND EVALUATION OF CONTROLLED POROSITY ORAL OSMOTIC PUMP TABLETS OF FUROSEMIDE
TEENA CHACKOCHEN THEKKAL, REBA RENJU, MANJU MARIA MATHEWS, BADMANABAN R Pages 110-113

[VIEW PDF](#)


FORMULATION AND EVALUATION OF TOPICAL GELS INCORPORATED WITH SOLID DISPERSIONS OF AN ANTIINFLAMMATORY DRUG
SETHU LEKSHMI, THERASE JOSE, MANJU MARIA MATHEWS, BADMANABAN R Pages 114-119

[VIEW PDF](#)

IN VITRO ANTI-BACTERIAL SCREENING OF DRYNARIYA QUERCIFOLIA

ASHNA T, LINS MARY JOY, SIYARA ANTONY, SINDU T J, SHEEBA MOL P, SHUJI T S, SOUMYA K GEORGE

Pages 120-123


 [VIEW PDF](#)

CASE REPORT(S)

GUILLAIN-BARRE SYNDROME: A PAEDIATRIC CASE SCENARIO IN A TERTIARY CARE HOSPITAL AT SOUTHERN INDIA

NEVIN JOSEPH, ALFIN BABY, ELDHOSE ELIAS GEORGE, GOPIKRISHNAN T.S, MERRIN JOSEPH

Pages 101-104

 [VIEW PDF](#)

[Announcements](#) || [Editorial Board](#) || [Indexing](#) || [Contact](#)

The publication is licensed under a [Creative Commons License \(CC BY-NC\)](#). [View Legal Code](#)

Copyright © 2023, JIAPSONline



Journal of Innovations in Applied Pharmaceutical Science [JIAPS]

Content available at: www.saap.org.in ISSN: 2455-5177



FORMULATION AND EVALUATION OF FLOATING CONTROLLED DRUG DELIVERY OF ANTI-ULCER DRUG LOADED MICROBALLOONS

Bindumol K C¹, Flowerlet Mathew^{*2}, Shalom Sunil¹, Angel Jose¹

¹Department of Pharmaceutics, Nirmala College of Pharmacy, Muvattupuzha

^{*2}Associate Professor, Department of Pharmaceutics, Nirmala College of Pharmacy, Muvattupuzha

Article History

Received: 06-10-2023

Revised: 26-10-2023

Accepted: 17-10-2023

Keywords: Floating microballoons, Famotidine, Eudragit RS100, HPMC K100M, Ethyl cellulose, Solvent evaporation method.

Abstract

An attempt has been made to develop floating drug delivery system for improving the drug bioavailability by prolongation of gastric residence time of Famotidine in the stomach. The floating microballoons were prepared using polymer Eudragit RS100, HPMC K100M and Ethyl Cellulose. Famotidine was used as the model drug. Nine formulations (F1 to F9) were prepared by varying the ratio of polymers. The prepared Famotidine loaded micro balloons were characterized for percentage yield, particle size analysis, surface morphology, micromeritic properties, drug entrapment efficiency, buoyancy studies and in-vitro drug release. The formulated micro balloons were free flowing. The micro balloons with Eudragit RS100 showed higher buoyancy when compared with HPMC K100M and Ethyl Cellulose. On the basis of micromeritic properties, particle size, percentage yield, morphology, buoyancy study, drug entrapment, optimum in-vitro drug release and satisfactory release kinetics, formulation F2 was selected as an optimum formulation.



This article is licensed under a Creative Commons Attribution-Non Commercial 4.0 International License.

Copyright © 2023 Author[s] retain the copyright of this article.



*Corresponding Author

Flowerlet Mathew

<https://doi.org/10.37022/jiaps.v8i3-S.532>

Production and Hosted by

www.saap.org.in

Introduction

Oral route is the most convenient and extensively used route for drug administration. This route has high patient acceptability, primarily due to easy of administration¹. Oral route of administration has been received more attention in the pharmaceutical field because of the more flexibility in the designing of dosage form than drug delivery design for other routes. Most of the oral controlled drug delivery systems rely on diffusion, dissolution or combination of both mechanisms, to release the drug in a controlled manner to the Gastro Intestinal Tract (GIT) and the drug profile data, such as dose, absorption properties and the quantity of drug needed, one can determine the desired release rate of the drug from controlled release dosage form

Drugs that are easily absorbed from the G.IT and having a short half-life are eliminated quickly from the blood circulation. To avoid this problem the oral controlled release formulations

have been developed, as these will release the drug slowly into the GIT and maintain a constant drug concentration in the serum for a longer period of time¹. More than 50% of drug delivery systems available in the market are oral drug delivery systems. These systems have the obvious advantages of ease of administration and patient acceptance. Ideal systems are not available. Thus, scientists try to develop systems that can be as close to an ideal system as possible. There are certain situations in which gastric retention is not desirable.

Gastroretentive dosage forms (GRDFs) are a drug delivery formulation that is designed to remain in the gastric region for a long period and significantly prolong the gastric retention time (GRT) of drugs [1] This technology has generated enormous attention over the last few decades owing to its potential application to improve the oral delivery of some important drugs. GRDFs prolonged retention in the upper gastrointestinal tract, greatly improve their oral bioavailability and/or their therapeutic outcome. Various techniques were used to encourage gastric retention of an oral dosage form. A significant approach for the treatment of gastric disorders can be achieved by floating drug delivery systems (FDDS)[2]. FDDS have a bulk density less than gastric fluids; they can float on the gastric juice in the stomach; the drug is released slowly at the

desired rate from the system [3]. A number of FDDS involving various technologies have been developed such as single and multiple-unit hydrodynamically balanced systems, single and multiple-unit gas generating systems, hollow microspheres, and raft forming systems

Hollow microspheres/microballoons are considered as one of the most promising buoyant systems [4]. They possess the unique advantages of multiple-unit systems as well as better floating properties because of central hollow space inside the microsphere. The drug release and better floating properties mainly depend on the type of polymer, plasticizer, and the solvents used for formulation. Commonly used polymers such as polycarbonate, cellulose acetate, calcium alginate, Eudragit S, agar and low methoxylated pectin were used in the preparation of hollow microspheres [5]. Hollow microspheres loaded with drugs in their outer polymer shell were prepared by simple solvent evaporation or solvent diffusion evaporation method to prolong the GRT of the dosage form. The microballoons floated continuously over the surface of an acidic dissolution media containing a surfactant for more than 12 hrs.

A peptic ulcer, also known as peptic ulcer disease (PUD), is a distinct breach in the mucosa of the stomach as a result of caustic effects of acid and pepsin in the lumen[4]. Histologically, peptic ulcer is identified as necrosis of the mucosa which produces lesions equal to or greater than 0.5 cm (1/5"). It is the most common ulcer of an area of the gastro intestinal tract that is usually acidic and thus extremely painful. Helicobacter pylori is one of the most common causes of peptic ulcer that colonizes the antral mucosa [5]. The immune system is unable to clear the infection, despite the appearance of antibodies. Thus, the bacterium can cause a chronic active gastritis (type B gastritis), resulting in a defect in the regulation of gastrin production by that part of the stomach, and gastrin secretion can either be increased, or as in most cases, decreased, resulting in hypo or achlorhydria[7]

Famotidine is a histamine H₂-receptor antagonist that inhibits stomach acid production, and it is commonly used in the treatment of peptic ulcer disease (PUD) and gastro esophageal reflux disease (GERD/GORD)[8]. Famotidine has no effect on the cytochrome P450 enzyme system, and does not appear to interact with other drugs. Famotidine binds competitively to H₂-receptors located on the basolateral membrane of the parietal cell, blocking histamine effects. This competitive inhibition results in reduced basal and nocturnal gastric acid secretion and a reduction in gastric volume, acidity and amount of gastric acid released in response to stimuli including food, caffeine, insulin, betazole, or pentagastrin.

Methods

Famotidine, Ethyl -cellulose, Eudragit RS100, HPMC K100M, Light liquid Paraffin, Acetone, Methanol, Dichloromethane, hydrochloric acid were the ingredients used in the formulation. Floating micro balloons were prepared by solvent evaporation method using 1000 mg of drug and different proportions of polymers were dissolved or dispersed in a mixture of dichloromethane and acetone (1:1) at room temperature. A solution of Famotidine was introduced to an external medium of light liquid paraffin (50 ml) in a 500 ml of beaker. The whole system was stirred at 1000 using

mechanical stirrer equipped with three blade propellers for 3-4 hours at 70.5°C to ensure the evaporation of the solvent. The solvent removal leads to polymer precipitation at o/w interphase of droplets forming cavity and thus mark them hollow to impart floating property. The prepared microballoons then filtered, washed with petroleum ether, air-dried and stored in desiccators.

Formula Code	Famotidine (mg)	Eudragit RS100 (mg)	HPMC K100M (mg)	Ethyl Cellulose (mg)	Dichloromethane: Acetone (ml)
F1(1:1)	1000	1000	-	-	1:1
F2(1:2)	1000	2000	-	-	1:1
F3(1:3)	1000	3000	-	-	1:1
F4(1:1)	1000	-	1000	-	1:1
F5(1:2)	1000	-	2000	-	1:1
F6(1:3)	1000	-	3000	-	1:1
F7(1:1)	1000	-	-	1000	1:1
F8(1:2)	1000	-	-	2000	1:1
F9(1:3)	1000	-	-	3000	1:1

Determination of Drug-Polymers Compatibility by FTIR Spectroscopy

FTIR spectra help to confirm the identity of the drug and to detect the interaction of the drug with the carriers. FTIR spectroscopy of pure drug and physical mixture of drug with polymers was carried out to check the compatibility between drug and polymers. The IR spectra of drug with polymers were compared with the standard IR spectrum of the pure drug.

Evaluation of Floating Microballoons

Micromeritic Studies

The prepared microballoons are characterized by their micromeritic properties, such as microballoon size, tapped density, Carr's compressibility index, Hausner's ratio and angle of repose.

Bulk Density

The bulk density is defined as the mass of powder divided by bulk volume. The bulk density was calculated by dividing the weight of the samples in grams by the final volume in cm³.

$$\text{Bulk density} = \frac{\text{Mass of microballoons}}{\text{Volume of microballoons before tapping}}$$

Tapped Density

Tapped density is the volume of powder determined by tapping by using a tapping cylinder containing weighed amount of sample. The cylinder containing own amount of microballoons was tapped for about 1 minute on a tapped density apparatus until it gives constant volume.

$$\text{Tapped density} = \frac{\text{Mass of microballoons}}{\text{Volume of microballoons after tapping}}$$

Carr's Compressibility Index

This is an important property in maintaining uniform weight.

$$\% \text{ Compressibility Index} = \frac{\text{Tapped density} - \text{Bulk density}}{\text{Tapped density}} \times 100$$

Hausner Ratio

A similar index like percentage compressibility index has been defined by Hausner. Values less than 1.25 indicate good flow, whereas greater than 1.25 indicates poor flow.

$$\text{Hausner's ratio} = \frac{\text{Tapped density}}{\text{Bulk density}}$$

Angle of Repose (θ)

It is essential that an accurate assessment of flow properties be made as early in the development process as possible so that an optimum formulation can be quickly identified. The angle of repose of each powder blend was determined by glass funnel method.

$$\tan \theta = h/r = \frac{\text{height of the pile}}{\text{radius of the powder cone}}$$

Percentage Yield

The measured weight of prepared microballoons was divided by the total amount of all the excipients and drug used in the preparation of the microballoons, which give the total percentage yield of floating microballoons.

$$\% \text{ Yield} = \frac{\text{Actual weight of product} \times 100}{\text{Total weight of excipients and drug}}$$

Particle Size Determination

Microballoon size was determined by using an optical microscope and the mean microballoon size was calculated by measuring 100 particles with the help of a calibrated ocular micrometer.

Morphological Study using SEM"

The morphological study was carried out by Scanning Electron Microscope. Carried out by using JEOL JSM-6390 a scanning microscope

Percentage Drug Entrapment

percentage Drug Entrapment was determined by taking weighed quantity of gisroballoons equivalent to 50 mg of the drug, thoroughly triturated and dispersed in so ml of 0.1 N HCl (pit-1.2) in 50 ml standard flask and stied for 24 hrs using a magnetic stirrer, filtered out and from the filtrate 1 mi was pipetted out and made up 10 10 ml with 0.1 N HCl and analysed spectrophotometrically at 265 nm against 0.1 N HI blank.

$$\% \text{ Drug Entrapment} = \frac{\text{Amount of drug actually present} \times 100}{\text{Theoretical drug load expected}}$$

In vitro Buoyancy Study™,

Microballoons equivalent to 40 mg of Famotidine were dispersed in a simulated gastric buffer pH 1.2 containing Tween 20 (0.02% w/v).The mixture was stirred with a paddle at 100 rpm. After 12 hrs, the floating microballoons and sinking particulate layer were separated by filtration. Particles of both types were dried at 40°C overnight in a desiccator until constant weight was achieved. weight was measured and buoyancy was determined

$$\text{Buoyancy \%} = \frac{W}{(W, + W_s)} \times 100$$

W, and Ws are the weight of the floating and settled microballoons

In-vitro Drug Release Study,

microballoons containing quantity equivalent to 40 mg of Famotidine by using US dissolution apparatus.

Result and Discussion

Determination of Drug-Polymers compatibility by FTIR Spectroscopy FT-IR spectra of the pure drug and the combination spectra of drug with the polymers, all the characteristic peaks of Famotidine were present in the

combination spectra as well thus indicating the compatibility of the drug with the polymers used.




Figure 24: FTIR Spectrum of Famotidine




Figure 25: FTIR Spectrum of Famotidine + Hydrogel M3100


Micromeritic Studies

The results of all formulations F1 to F9 of Floating microballoons are shown

Formulation code	Bulk density (g/cc)	Tap density (g/cc)	Carr's Index	Hausner ratio	Angle of repose()
F1	0.425	0.451	5.7650	1.0612	26.27
F2	0.577	0.624	7.5321	1.0815	27.88
F3	0.723	0.772	6.3472	1.0678	29.01
F4	0.372	0.392	5.1020	1.0538	28.34
F5	0.548	0.580	5.5172	1.0584	29.29
F6	0.631	0.673	6.2407	1.0666	30.03
F7	0.728	0.772	5.6994	1.0604	26.37
F8	0.479	0.510	6.0784	1.0647	27.76
F9	0.296	0.316	6.3291	1.0676	28.13


Percentage Yield

The percentage of production yield of the prepared floating microballoons of Famotidine was in the range of 67-98.25% being the highest for formulation F6 and lowest for formulation F7.




In-vitro Buoyancy Study

The Percentage buoyancy of the prepared floating microballoons of Famotidine was in the range of 57.64-72.01%, being the highest for formulation F2 and lowest for formulation F4





Particle Size Determiation

The Mean Particle Size of the prepared floating microballoons of Famotidine was in the range of 52.11-64.21 um being the highest for formulation F3 Famotidine lowest for formulation F7




Morphological Study using SEM

Particle shape and surface morphology had been evaluated using SEM and the prepared microballoons were found to be almost spherical in nature and possessing more or less smooth outer surface.




Percentage Drug Entrapment

The Percentage Drug Entrapment of the prepared floating microballoons of Famotidine was in the range of 70.0540-91.7496%, being the highest for formulation F3 owest for formulation F7.



In-vitro Drug Release Study

In-vitro release of Famotidine from microballoons was determined by carrying out dissolution test using USP basket method at a stirring rate of 100 pm at temperature. The in-vitro drug release of prepared formulations was found to be sustained in a period of 12 hours and it was highest for formulations containing Budragit R\$100(1:2) and lowest for formulations containing HPMC K100M (1:3). The cumulative release of Famotidine significantly decreases with increasing polymer concentration.



Kinetics

The in-vitro drug release data of F2 was applied to various kinetic models like zero order kinetics, first order kinetics, Higuchi plot, Korsmeyer-Peppas plot and Hixson-Crowell plot to predict the drug release kinetics mechanism. The

formulation F2 was best fitted with Higuchi model and it was found to be following Fickian diffusion as release mechanism.



Discussion

The percentage yield obtained in all the formulations was good and in the range of 67-98.25% w/w. The percentage drug entrapment of all the formulations was found good and in the range of 70.0540-91.7496% w/w.[9] The particle size analysis indicated that the size of microballoons was found to increase with increase in the concentration of polymer and range of particle size of the prepared microballoons was found to be 52.11-64.21. The SEM photograph of microballoons revealed that the prepared microballoons were almost spherical with a more or less smooth outer surface. Further studies are required to establish the in-vivo bioavailability and in-vitro & in-vivo correlation of bioavailability of the microballoon formulations prepared [10]. This can be used for the further studies of floating microballoons.

Conclusion

Floating is one of the important approaches to the gastric retention of the dosageform. Famotidine with a low oral bioavailability of 40-45% and half-life of 2.5-3.5hours is an example of drugs that could benefit from increased residence in the stomach.

Based on in-vitro buoyancy study and in-vitro drug release characteristics, the formulation F2 was selected as the best formulation, as it showed a percentage buoyancy of 72.01% and a higher in-vitro drug release profile[11] (88.8370% of drug released in 12 hours which was sustained up to 12hours.The in-vitro drug release data of F2 was applied to various kinetic models like zero order kinetics, first order kinetics, Higuchi plot, Korsmeyer-Peppas plot and Hixson-Crowell plot to predict the drug release kinetics mechanism[13].Theformulation F2 was best fitted with Higuchi model and it was found to be following Fickian diffusion as release mechanism.Future scope of floating drug delivery systems lies in the modification techniques of polymer carriers to improve their floating property and to make them more target-specific.Flow properties, suggesting that, in future they could be easily and successfully packed and developed into a capsule dosage form.

The prepared microballoons proved to be a potential candidate as a microparticulate controlled release drug delivery device in this era of patenting novel and controlled release formulations.

Acknowledgement

Not declared

Funding

No

Conflict of interest

No Conflict of interest

Ethical approval and Inform Consent

Not Required

Reference

- Gupta S, Mali RR, Goel V. Novel study in fast dissolving drug delivery system: A review. *Indian J Pharm Biol Res* 2015;3(1):93-107.
- Nayak AK, Maji R, Das B. Gastroretentive drug delivery systems: A review. *Asian J Pharm Clin Res* ;3(1):1-10.
- Kiss D, Zelkó R. Gastroretentive dosage forms. *Acta Pharm Hung* ;75(3):169-76
- Beers MH, Fletcher AJ, Jones T, Porter R, Berkwitz M, Kaplan J. *The Merck manual of medical information*. 2nd ed. New York: Pocket Books; Merck manual of medical information. 2nd ed. New York: Pocket Books; 2003;390(10094):613-624
- Narayanan M, Reddy KM, Marsicano E. Peptic Ulcer isease and Helicobacter pylori infection. *Mo Med*. 2018 May-Jun;115(3):219-224.
- Lanas A, Chan FKL. Peptic ulcer disease. *Lancet*. 2017 Aug 05 Beers MH, Fletcher AJ, Jones T, Porter R, Berkwitz M, Kaplan J.
- K. Lucija, J. Jelena, S. Robert, R.L. Nikola, V. Aleksandar. Peptic Ulcer Disease: A Brief Review of Conventional Therapy and Herbal Treatment Options. *J Clin Med*. 2019; 8(2): 179.
- Lakshmanamurthy G, JeevanandhamS, Kumar GP. Formulation and in-vitro evaluation of gastro retentive Famotidine hollow microspheres. *Journal of Drug Delivery & Therapeutics* 2013;3(6):207-14.
- Deepaa MK, Karthikeyan M. Cefpodoxime proxetil floating microspheres: Formulation and in-vitro evaluation. *Iranian Journal of Pharmaceutical Sciences* 2009;5(2):69-72.
- Nithyashree RS. Kumar S. Gosh T. Floating Microballoon-A Novel Formulation For Gastrointestinal Diseases, *Journal of Pharmaceutical Hajare P Rachh P R. Gastroretentive microballoons: a novel approach for drug delivery. International Journal of Pharmaceutical Sciences and Research*. 2020;11(3):1075-83
- Sciences and Research. Bandana S, Jovita K, Manisha P and Sarag SA. Formulation and evaluation of floating microspheres of famotidine. *International Journal of Pharm Tech Research* 2010 April-June; 2(2):1415-20020;12(1):74-78; 0975-1459.
- Maryam K and Ali B. Preparation and in-vitro evaluation of a microballoon delivery system for Theophylline. *Iranian Journal of Pharmaceutical Research*
- K.D. Tripathi. *Essentials of medical pharmacology*, 8th edition, Jaypee Brothers Medical publication (P) Ltd: New Delhi, 2019; 698 p.
- R. Bairam, H. Kumar Tatapudi, N. Gajji, S. Harun Rasheed. UV Spectrophotometric and HPLC Method for Quantification of Ticagrelor in Bulk and Tablet Dosage

Form. Asian Journal of Pharmaceutical Analysis. 2022; 12(3):159-5.

15. Benoit D.S., Overby C.T., Sims K.R., Jr., Ackun-Farmmer M.A. Biomaterials Science. Elsevier; Amsterdam, The Netherlands: 2020. Drug delivery systems; pp. 1237–1266.