



NIRMALA COLLEGE OF PHARMACY MUVATTUPUZHA



NATIONAL CONFERENCE

Nano-based Drug Delivery Systems; Recent Developments and Future Prospects

7 OCTOBER 2023

ASSOCIATING PARTNERS

INDIAN PHARMACEUTICAL ASSOCIATION



**JOURNAL OF INNOVATIONS IN
APPLIED PHARMACEUTICAL SCIENCES**



**INNOVATION AND
ENTREPRENEURSHIP
DEVELOPMENT CENTRE**



**INSTITUTION'S
INNOVATION
COUNCIL**
(Ministry of HRD Initiative)

EFFICIENT MICROWAVE SYNTHESIS OF COUMARIN DERIVATIVES WITH EVALUATION OF THEIR ANTIOXIDANT AND ANTI-INFLAMMATORY PROPERTIES
ANZIYA P A, SARANYA T S, ANJALI K, ANJALI KRISHNA, SINI BABY, DIVINE P DANIEL Pages 124-130

[VIEW PDF](#)

COSMETIC USE RELATED ADVERSE EVENTS AND NEED FOR COSMETOVIGILANCE
MERRIN JOSEPH, KARISHMA SHAJI, MAHIN T M, NANDANA P B, KRISHNA DAS Pages 64-71

[VIEW PDF](#)

A RETROSPECTIVE STUDY OF CLINICAL PROFILE OF VIPER BITE CASES IN SELECTED HOSPITALS IN CENTRAL KERALA
ANUMOL SAJU, ANTRIYA ANNIE TOM, ABY PAUL, SWAPNA SAJU, DONA JOHNSON, JESYLN JOE THOMAS, KUTTIKADEN Pages 72-74
JOY STEFFI, JOYAL M JOLL

[VIEW PDF](#)

FORMULATION AND EVALUATION OF HERBAL TOOTHPASTE CONTAINING EUPATORIUM TRIPLINERVISLEAF EXTRACT
VIDYA PETER, ROSNA BABU , SHERRY SEBASTIAN, ANGEL JAEMON, ANGEL JAEMON, ANAGHA V T, JEEVAN SAJEEV Pages 36-41

[VIEW PDF](#)

IN VITRO SCREENING OF ICACINACEOUS PLANTS INDIGENOUS TO KERALA
DR.ELIZABETH ABRAHAM P, FRINTO FRANCIS, PRADEEP R NAIR, ATHUL RAJ, RAJI RAJAN, ANAMIKA K. NAIR, PROF.DR.BADMANABAN.R Pages 51-58

[VIEW PDF](#)

FORMULATION AND EVALUATION OF BUCCAL FILM OF AN ANTIHYPERTENSIVE DRUG
ASHINAA BENEDICT, IRIN ROSE PAUL, DR. MANJU MARIA MATHEWS, DR. BADMANABAN R Pages 75-80

[VIEW PDF](#)

A PROSPECTIVE SURVEY TO ASCERTAIN THE SYMPTOMS, HEALTH ISSUES AND SUBSEQUENT OTC MEDICATION USAGE DURING MENSTRUATION AMONG COLLEGE STUDENTS
MINTU GEORGE, ANAGHA MELBIN, MARY PAUL DOMINIC, RESHMA DOMINIC, AYSHA SAJA P.S, JOBIN KUNJUMON Pages 81-84
VILAPURATHU

[VIEW PDF](#)

A CROSS SECTIONAL STUDY TO ANALYSE THE ADR REPORTED IN A HOSPITAL DURING THE PAST THREE YEARS
SANGEETHA SUKUMARAN, VARSHA ELIZABETH JOBY, AMALA JOSEPH, APARNA JESTIN, JITHIN N P, SUMAYYA B Pages 85-89
MUHAMMED, SUNU SEBASTIAN, JOBIN KUNJUMON VILAPURATHU

[VIEW PDF](#)

FORMULATION AND EVALUATION OF PREUNGUAL DELIVERY SYSTEM CONTAINING EUGENOL FOR THE TREATMENT OF ONYCHOMYCOSIS
MINI ELIAS, FLOWERLET MATHEW, GOURISREE T, ANILA RAJAN, ASHLY DAVIS Pages 90-94

[VIEW PDF](#)

FORMULATION AND EVALUATION OF FLOATING CONTROLLED DRUG DELIVERY OF ANTI-ULCER DRUG LOADED MICROBALLOONS
BINDUMOL K C, FLOWERLET MATHEW, SHALOM SUNIL, ANGEL JOSE Pages 95-100

[VIEW PDF](#)

PREPARATION AND EVALUATION OF FLOATING DRUG DELIVERY SYSTEM (FDDS) CONTAINING AN ANTIVIRAL DRUG
TEENA MOHAN, MARIYA SUNNY, MANJU MARIA MATHEWS, BADMANABAN R Pages 105-109

[VIEW PDF](#)

FORMULATION AND EVALUATION OF CONTROLLED POROSITY ORAL OSMOTIC PUMP TABLETS OF FUROSEMIDE
TEENA CHACKOCHEN THEKKAL, REBA RENJU, MANJU MARIA MATHEWS, BADMANABAN R Pages 110-113

[VIEW PDF](#)

FORMULATION AND EVALUATION OF TOPICAL GELS INCORPORATED WITH SOLID DISPERSIONS OF AN ANTIINFLAMMATORY DRUG
SETHU LEKSHMI, THERASE JOSE, MANJU MARIA MATHEWS, BADMANABAN R Pages 114-119

[VIEW PDF](#)



IN VITRO ANTI-BACTERIAL SCREENING OF DRYNARIYA QUERCIFOLIA

ASHNA T, LINS MARY JOY, SIYARA ANTONY, SINDU T J, SHEEBA MOL P, SHUJI T S, SOUMYA K GEORGE

Pages 120-123

[VIEW PDF](#)

CASE REPORT(S)

GUILLAIN-BARRE SYNDROME: A PAEDIATRIC CASE SCENARIO IN A TERTIARY CARE HOSPITAL AT SOUTHERN INDIA

NEVIN JOSEPH, ALFIN BABY, ELDBOSE ELIAS GEORGE, GOPIKRISHNAN T.S, MERRIN JOSEPH

Pages 101-104

[VIEW PDF](#)

[Announcements](#) || [Editorial Board](#) || [Indexing](#) || [Contact](#)

The publication is licensed under a [Creative Commons License \(CC BY-NC\)](#). [View Legal Code](#)

Copyright © 2023, JIAPSONline





Journal of Innovations in Applied Pharmaceutical Science [JIAPS]

Content available at: www.saap.org.in ISSN: 2455-5177



IN VITRO ANTI-BACTERIAL SCREENING OF *DRYNARIYA QUERCIFOLIA*

Ashna T¹, Lins Mary Joy*², Siyara Antony¹, Sindu T J¹, Sheeba Mol P¹, Shiji T S¹, Soumya K George¹

¹ Nirmala College of Pharmacy, Muvattupuzha

² Associate Professor, Department of Pharmacognosy, Nirmala College of Pharmacy, Muvattupuzha

Article History

Received: 06-10-2023

Revised: 28-10-2023

Accepted: 19-10-2023

Keywords: *Drynaria quercifolia*, phytoconstituent, anti-bacterial, extract.



Abstract

Purpose: To make scientific validation of antibacterial activity of ethnomedical fern *Drynaria quercifolia*. **Method and results:** The rhizome extracts were prepared using different solvents like benzene, ethyl acetate, chloroform, ethanol & water. The percentage yield was 7, 3, 2, 3, 3.1 & 6.4 respectively. Preliminary phytochemical screening revealed the presence of maximum constituents in the ethyl acetate extract. The anti-bacterial activity was screened against both gram-positive and gram-negative bacteria, and the zone of inhibition was measured. The results were compared with standard Ampicillin (250µgm) and ethyl acetate was found to be most effective against *Bacillus subtilis*. **Conclusion:** In the antibacterial study ethyl acetate extract showed good antibacterial activity even in microgram quantities and is found to be very effective against *Bacillus subtilis*. Based on the studies it can be concluded that *Drynaria quercifolia* can be effectively used as an anti-bacterial agent.

This article is licensed under a Creative Commons Attribution-Non Commercial 4.0 International License. Copyright © 2023 Author[s] retain the copyright of this article.



*Corresponding Author

Lins Mary Joy

<https://doi.org/10.37022/jiaps.v8i3-S.537>

Production and Hosted by

www.saap.org.in

Introduction

Drynaria is an epiphytic or pterophilous fern widely distributed throughout Asia to North-East Australia and Africa. Four species are reported to occur in India. *Drynaria quercifolia* alone is distributed in South India, in the plains or very low down in mountains, on trees or rocks.

The rhizomes are short thick fleshy creeping, up to 3 cm in diameter, adpressed to the substratum tightly, densely clothed with paleae all over. Paleae linear- lanceolate or lanceolate, peltate with a short stalk, red brown, non-clathrate, gland-tipped, gland deciduous, margins dentate-ciliate.

The fronds are seasonal, dimorphic, coriaceous or sub coriaceous of two types; The sterile fronds become brown on ageing and are small and somewhat concave. They vary in size from 7.5 to 30 cm (l) and 18-20 cm(w), cordate oval, lobate - pinnatifid sessile, lateral veins prominent and closely covering rhizome.

Taxonomical Classification	Regional Names/ Vernacular names
Domain: Eukaryote Kingdom: Plantae Sub kingdom: <i>Viridaeplantae</i> Phylum: <i>Tracheophyta</i> Sub phylum: <i>Pteridophytina</i> Infraphylum: <i>Filices</i> Class: <i>Filicopsida</i> Order: Filicales Family: <i>Polypodiaceae</i> Sub family: <i>Vaccinioideae</i> Tribe: <i>Androcdeae</i> Botanical name: <i>Drynaria quercifolia</i> Synonym: <i>Polypodium quercifolium</i>	Hin: Asvakatri, Katikapan Mal: Matilpanna, Pannakkilannu San: Asvakatri, Tam: Pannakkilanku



Image 1: *Drynaria quercifolia*



Image 2: *Drynaria quercifolia* rhizome

Medicinal uses

Anti-bacterial, Anti-fungal, Antipyretic, Anthelmintic, Anti-inflammatory, Hepatoprotective, Antiulcer, Anticancer, Anti-diabetic etc.

Chemical constituents

Phytoconstituents isolated from the plant include friedelin, epifriedelinol, β -sitosterol, β -amyrin, 3,4-dihydroxybenzoic acid, β -sitosterol 3- β -D-glucopyranoside, naringin, naringenin and acetyl lupeol.

Materials and methods

Analytical parameters

Determination of total ash values

2 gm of rhizome powder was separately weighed and taken in a pre-weighed silica crucible. The powder was incinerated at a temp above 450°C until free from carbon and then cooled in a desiccator. The weight of total ash was taken and the percentage yield was calculated with reference to air dried sample.

Determination of Acid insoluble ash value

The total ash obtained was boiled with 25ml of 2 N HCl for 5 minutes. The insoluble matter was collected on an ashless filter paper and washed with hot water until it is neutral. The filter paper is then incinerated in a crucible and cooled in a desiccator and weighed. Percentage of acid insoluble ash with reference to air dried drug was calculated.

Determination of Water-soluble ash value

The total ash obtained was boiled with 25 ml of water for 5 minutes. The insoluble matter was collected on an ashless filter paper and washed with hot water until it is neutral. The filter paper is then incinerated in a crucible and cooled in a desiccator and weighed. Percentage of acid insoluble ash with reference to air dried drug was calculated.

Sl.no	Ash Value	Percentage ash (w/w)
1.	Total ash value	8%
2.	Acid insoluble ash value	12.5%
3.	Water soluble ash value	50%

Determination of extractive values

Determination of water-soluble extractive value

5gm of the coarsely powdered drug was accurately weighed and macerated with 100 ml of chloroform water in a stoppered flask for 24 hours shaking frequently for 6 hours and allowing to stand for 18 hours. The mixture was filtered and 25 ml of the filtrate was evaporated to dryness in a flat-bottomed shallow dish, dried at 105°C, cooled in desiccator and weighed. The percentage w/w of water-soluble extract with reference to the air-dried drug was calculated.

Determination of alcohol soluble extractive value

This was determined by the method described under water-soluble extractive value using 100 ml of 90% alcohol instead of chloroform water taking adequate precautions to avoid loss of alcohol during filtration.

Nature of extract	Percentage (w/w)
Water soluble extractive value	2.2%
Alcohol soluble extractive value	1.4%

Extraction procedures

Collection & Processing of *Drynaria quercifolia*

The rhizomes are collected from Idukki and identified in botany department of Nirmala College, Muvattupuzha. They are cleaned properly, cut into pieces and dried in shade. The fine powder prepared by blender is used for extraction.

Method of extraction

31g were accurately weighed & extracted in Soxhlet apparatus using solvents benzene (2.7), ethyl acetate (4.3), chloroform (4.4), ethanol (5.2) and water (9) in the increasing order of polarity. Each time before extracting with the next solvent the powdered material was dried.

Percentage Yield of *Drynaria quercifolia*

Sl. No	Extracts	Weight in grams	%yield (w/w)
1.	Benzene	2.17g	7%
2.	Ethyl acetate	0.971g	3%
3.	Chloroform	0.713g	2.3%
4.	Ethanol	0.991g	3.1%
5.	Water	2.01g	6.4%

Preliminary Phytochemical Analysis of *Drynaria quercifolia* extracts

All extracts of *Drynaria quercifolia* rhizome were subjected to qualitative chemical tests for the identification of various phytoconstituents.

	Benzene	Ethyl acetate	Chloroform	Ethanol	Water
Steroids	-	+	-	-	-
Alkaloids	-	-	+	-	-
Glycosides	-	+	-	-	-
Flavonoids	-	+	-	-	-
Carbohydrates	+	+	+	+	-
Fats&oils	+	+	+	+	-
Proteins	-	-	+	+	+
Saponins	+	+	-	-	-

Identification of Active Constituents by Thin Layer Chromatography Procedure

The benzene, ethyl acetate, chloroform, ethanol, water was used to dissolve the samples.

Solvent system: 15% methanol in chloroform

Detection technique: -U. V

The TLC plates (7.6 x 2.5cm) were prepared with silicagel G. A slurry of adsorbent was prepared by triturating it with water in a mortar. The slurry was then spread uniformly over the glass plates and thin layer was allowed to set. The plates after spreading were air dried and then activated by heating in an oven at 100-110°C for 30-60 minutes. Then the various extracts were deposited as a spot with the help of capillary tubes. After drying the chromatogram were developed in solvent system Methanol: Chloroform (15:85, vol: vol) by placing in a developing tank which was paper lined so that the atmosphere inside was saturated with solvent phase. The plates were removed when the solvent front reaches 3/4 of the plate. The solvent front was marked and the plates were allowed to dry. The spots were detected by UV.

The extracts were separated into different spots. The Rf values were calculated by measuring the distance travelled by the solute and the solvent.

$$Rf \text{ Value} = \frac{\text{Distance travelled by solute from the origin}}{\text{Distance travelled by solvent front from the origin}}$$

Rf Value of Different Compounds Isolated from *Drynaria quercifolia* Rhizome

Sl. No.	Name of the extract	Distance travelled by solute in cm	Colour of the spot	Number of spots	Distance travelled by solvent front in cm	Rf value
1.	Benzene	4.2	Yellow	1	5.4	0.77
2.	Chloroform	4	Yellowish green	1	5.5	0.72
3.	Ethyl acetate	1.5 3 4.3	Yellow Blue blue	3	5.7	0.26 0.52 0.75

Evaluation of Antibacterial Activity

Preparation of Plant Extract

Various extract of *Drynaria quercifolia* prepared with the help of Soxhlet apparatus. 250 mcg/ml solutions were prepared by dissolving the various extracts in the corresponding solvents.

Preparation of Media

Each ingredient (Beef extract-10gm, peptone -10gm, sodium chloride- 5gm) except agar was dissolved in the appropriate volume of distilled water. The pH of the fluid medium is determined with a pH meter and adjusted to pH 7 by using 1N HCl or 1N NaOH. Agar powder was added and the medium was heated to dissolve the agar to form a clear liquid. The medium was dispensed into flasks, plug the flasks containing medium by using nonadsorbent cotton. The media was sterilized at 121° C at 15 lbs pressure for 15 mins in an autoclave. Allow the flask to cool up to 50°C and pour the medium quickly into sterile petriplates (sterilized by hot air oven by heating at 160°C for hr) under aseptic conditions. Allow the medium to cool and to produce solid agar plates.


Procedure

Invitro antibacterial screening was carried out against two Gram positive. [*Bacillus subtilis*, *Staphylococcus aureus*] and two Gram negative [*Escherichia coli*, *Pseudomonas aeruginosa*] bacteria.


Disc diffusion method, was used for antibacterial screening, 20 ml quantities of nutrient agar were plated in petridish with 0.1 ml of each bacterial culture. Filter paper discs (6 mm in diameter) impregnated with respective samples [Benzene, Ethyl acetate, Chloroform, Ethanol and Water fraction] were placed on the test organism seeded plates. Benzene, Ethyl acetate, Chloroform, Ethanol and Water were used to dissolve the samples and were completely evaporated before application on the test organism seeded plates. Blank disc impregnated with

solvents Benzene, Ethyl acetate, Chloroform, Ethanol, Water followed by drying off were used as negative control. The activity was determined after 18 hrs of incubation. The zone of inhibition produced by samples was then compared with the standard antibiotic (Ampicillin).

Percentage inhibition of *Bacillus subtilis*, *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa* on ethyl acetate extracts of *Drynaria quercifolia*.



Antibacterial Activity of Ethyl Acetate Extract of Rhizomes of *Drynaria quercifolia* Against *Bacillus subtilis*.



Results and Discussion

The percentage yield of Benzene, Ethyl acetate, Chloroform, Ethanol, Water extracts of *Drynaria quercifolia* were found to be 7, 3, 2.3.3.1 and 6.4 respectively. Analytical parameters like Ash value and Extractive values were determined as a mode of identification of the fern. The presence of maximum active constituents like steroids, glycosides, flavonoids, carbohydrates, fats and saponins in ethyl acetate extract was confirmed by phytochemical screening and is decided to scientifically validate its antibacterial activity. The TLC of

Benzene, Ethyl acetate and Chloroform extracts were performed using solvent system, Methanol: Chloroform (15:85). The three different spots were detected using UV light and their Rf values were 0.75, 0.52 and 0.26 respectively. The antibacterial activity of the ethyl acetate extract of *Drynaria quercifolia* showed a percentage activity of 89 %, 75 %, 60 % & 56 % against *B. subtilis*, *E. coli*, *S. aureus* & *P. aeruginosa* respectively.

Conclusion

The antibacterial study ethyl acetate extract shows good antibacterial activity even in microgram quantities and is found to be very effective against *Bacillus subtilis*. So *Drynaria quercifolia* can be used effectively as an anti-bacterial agent.

Acknowledgement

We express our sincere gratitude to the management of Nirmala College of Pharmacy Muvattupuzha, Kerala, for providing facilities for the completion of the work. Also we extend heartfelt thanks to Botany Department, Nirmala College, Muvattupuzha for identification of the plant identification of the plant.

Conflict of interest

Authors are declared that no conflict of interest.

Funding

Funded by Nirmala College of Pharmacy, Muvattupuzha

Reference

1. H.N. Srivastava. Fundamentals of pteridophytes. 13th ed., Pradeep publications, 2002,1-16.
2. Santhosh Nampy, P.V. Madhusoodanan. Fern flora of South India. Daya Publishing house, 1998; 145-49.
3. Kirtikar KR, Basu BD. Indian medicinal plants. 4 Vol., 2nd ed., Dehra Dun, India, Dehra Dun Publisher Ltd. 1994;
4. Indian medicinal plants. 2nd Vol., 345.
5. Ramesh.N. Viswanathan M.B., Saraswathy A., Balakrishnan.K. Et.al. Phytochemical and antimicrobial studies on *Drynaria quercifolia*.
6. Khan. A., Haque.E., Mukhlesure et.al. Isolation of antibacterial constituents from rhizome of *Drynaria quercifolia* and its sub-acute toxicological studies.
7. Alam Khan, Ekramul Haque.et.al. Antipyretic activity of rhizome of *Drynaria quercifolia*.
8. M. Kandhasamy, K.D. Arunacahalam et.al. *Drynaria quercifolia* (L.) J. Sm: A potential resource for antibacterial activity.
9. Liu, HC: Chen, RM: Jain, WC: Lin, YL. Cytotoxic and antioxidant effects of the water extract of the traditional Chinese herb gusuibu (*Drynaria*) on rat osteoblasts.
10. www.google.com
11. C.K. Kokate, A.P. Purohit.et.al Text book of Pharmacognosy. 593-597, 112-117.
12. Dr. A.V. Kasturi, Dr. K.R. Mahadi.et.al Pharmaceutical analysis. 2 Vol., 23-26, 251.
13. A.K. Aswathy, G.S. Bisht: et.al Invitro antimicrobial activity of Swertiachirata Chirata. Indian j. Nat. Prod. 21(2), 27, 27-29.
14. Natural product Radiance - a bimonthly Journal of natural products July-August 2006 5th Vol. 257-344.
15. Pulok.K. Mukherjee. Quality control of herbal drugs. 446-447.
16. Harbone.T.B. Phytochemical method (A guide to modern technique of plant analysis). I edition, 10-12.
17. Bentlys Text book of Pharmaceutics.8thed., Published by All India. Traveller Book Seller,2005;173,181
18. Ananthanarayan and Panikers Text Book of Microbiology.7th ed., Orient Longman Private Ltd.34
19. Quality control methods for medicinal plant material, World health Organisation, Geneva.A.I.T.B. S Publishers and Distributers (Regd) 2002,22,28,29
20. Indian Pharmacopoeia 1996, Vol-11;Govt. of India, Ministry of Health and Family Welfare,53,54
21. Trease and Evans, Pharmacognosy.15ed.,138
22. Thin-Layer Chromatography, Springer International Edition. 2nd ed.,
23. G Prasanna, R Anuradha, A Comprehensive Review on Phytopharmacological Activitiesof *Drynaria quercifolia*, International Journal of Pharmacognosy and Phytochemical Research 2016; 8(8); 1304-1313.
24. Subash Kumar Gupta, Dr.Jnan Bikash Bhandarietal, International Journal of Creative Research Thoughts (IJCRT), Volume 9, Issue 4 April 2021,5464-5473.
25. Anuja GI, Latha PG, Shine VJ, Suja SR,etal., Anti-inflammatory and analgesic properties of *Drynaria quercifolia* (L.) J. Smith,J,Ethnopharmacology 2010 Nov 11;132(2):456-60
26. Mudavath Ravi Naik , Praveen Thaggikuppe Krishnamurthy,A Review on Phytochemicals and Pharmacological Activities of the medicinal plant *Drynaria quercifolia*. European Chemical Bulletin, Volume 12 Issue 8(2023).
27. Prasanna Getal , In Vitro Evaluation Of Antidiabetic And Cytotoxicity Potentials Of The Rhizome Extract Of *Drynaria Quercifolia* (L.) J. Smith, Vol 12 Issue 11 November 2019.
28. Alam Khanetal., Antipyretic Activity of Rhizome of *Drynaria quercifolia*. in Rabbit, Pharmaceutical Biology, Issue 4 ,Volume 45, 2007 ,Pages 312-315.
29. Lakshman RRetal.Screening the anticancer activity, cytotoxicity and anti-osteoarthritic activity of *Drynaria quercifolia* through Saos2 cell line studies (2023) Volume 11, Issue 2.
30. Pradeep Kambojetal., Hepatoprotective Effect of *Drynaria quercifolia* Fronds Hydroalcoholic Extract and Isolated Constituent against CCl4-Induced Hepatocellular Damage,Journal of Pharmaceutical Research International, 2013 - Volume 3 [Issue 4].

